

Use the following posts for your clinic's social media to raise awareness about Intestinal Parasites.

Roundworms are the most common parasitic worms found to infect dogs. In fact, it has been found that almost ALL dogs are infected with roundworms at some point in their lives.

Did you know?

Many dogs won't show initial signs of a roundworm infection. If the infections continue, untreated, it worsens your dogs may experience diarrhea, vomiting weight loss.

FAQ: Can people be harmed by roundworms?

A: Roundworm eggs can accumulate in soil where your pet does his "business". These worms can be transferred to you or your family members from infected soil or your pet's feces. Once infected, these worms pose a significant health risk, especially in children, where they can cause eye, lung and heart damage.

Unfortunately, it is fairly common for a dog to become infected with an internal or external parasite during his or her lifetime. These pesky parasites can affect your loved one in a variety of ways, sometimes even causing serious health problems. Fortunately, we can help! Call our office today and we can find the prevention protocol that fits your pet and family's lifestyles best.

Did you know?

Dogs can pick up hookworms simply though skin contact with infected soil or through ingestion of the dirt through routine cleaning.

Whipworms are one of the four most common intestinal parasites that can infect your dog. Dogs become infected with whipworms, as most other intestinal parasites, through infected soil. Because whipworms can be more difficult to diagnosis than other intestinal parasites, it is important your dog doesn't miss his annual test and monthly worming medication.